

TABLE 2. 1998 SISKIYOU INTERMOUNTAIN WINTER BARLEY TEST

ENTRY	YIELD LBS/ACRE	TEST WT (LBS/BU)	1000 KERNEL		PLANT HT (IN)	LODGING		STRIPE RUST
			WT (GMS)	SHATTER		AT SOFT DOUGH	LODGING AT HARVEST	
204 STEPTOE	3110 (09)	45.9	39.7	1.5	50	7.0	7.3	8.0
567 SCIO	4520 (03)	48.8	40.8	1.0	43	1.3	1.3	3.0
568 HESK	3920 (07)	47.1	36.2	1.0	44	2.3	2.3	3.8
574 BOYER	4180 (06)	48.9	38.4	1.0	46	1.5	1.3	2.3
697 EIGHT TWELVE	4270 (05)	46.5	34.9	1.0	42	1.0	3.0	4.3
748 WESTBRED SPRINTER	4450 (04)	51.4	39.9	1.0	45	1.8	1.5	1.0
878 SHOWIN	2950 (10)	43.1	31.6	1.0	40	1.0	3.3	4.5
908 KOLD	5570 (01)	48.9	36.9	1.0	45	1.5	2.3	1.0
950 SUNSTAR DOUBLE	3770 (08)	47.5	32.0	1.0	45	1.0	1.3	7.5
952 SUNSTAR PRIDE	1000 (12)	41.4	26.1	1.3	41	1.5	2.5	8.0
964 STRIDER	5490 (02)	48.4	39.2	1.0	44	1.3	1.8	1.5
982 88Ab536	2920 (11)	48.5	35.8	1.8	42	4.5	6.3	3.8
MEAN	3850	48.2	36	1.1	44	2.1	2.8	4
CV	12.6	1.3	6.5	22.3	6	45.8	40.4	26.4
LSD (.05)	700	1.1	5.1	0.4		1.4	1.6	1.5

Rating scale for diseases (area of flag-1 leaf affected at soft dough stage), lodging and shatter: 1=0-3%, 2=4-14%, 3=15-29%, 4=30-49%, 5=50-69%, 6=70-84%, 7=85-95%, 8=96-100%.

Numbers in parentheses indicate relative rank in column.