

TABLE 3. 2000 SISKIYOU INTERMOUNTAIN WINTER BARLEY TEST

Entry	Name	Test 1000			Plant				
		Yield (lbs/acre)	Wt (lbs/bu)	Kernel Wt (g)	Shatter	Ht (in)	Lodging (Soft dough)	Lodging (Harvest)	Stripe Rust
204	STEPTOE	5160 (08)	50.8	49.7	2.5	47	2.5	3.3	6.8
567	SCIO	5770 (07)	49.9	42.1	1.8	42	4.8	5.0	1.0
568	HESK	5960 (06)	47.0	37.4	1.0	44	8.0	8.0	1.0
574	BOYER	6300 (05)	45.2	35.6	1.5	45	7.5	8.0	1.0
697	EIGHT TWELVE	6640 (03)	50.3	41.4	1.5	41	2.8	5.0	1.0
748	WESTBRED SPRINTER	6730 (02)	47.9	39.0	1.0	42	7.0	7.3	1.0
908	KOLD	7240 (01)	48.4	36.1	1.0	42	8.0	8.0	1.0
964	STRIDER	6470 (04)	50.8	42.1	1.3	44	8.0	7.8	1.0
MEAN		6280	48.8	40.4	1.4	43	6.1	6.5	1.7
CV		19.9	2.3	1.8	53.4	2.9	20	16.9	10.3
LSD (.05)			2.6	1.8		3.0	1.8	1.6	0.3

Rating scale for diseases (area of flag-1 leaf affected at soft dough stage), lodging and shatter: 1=0-3%, 2=4-14%, 3=15-29%, 4=30-49%, 5=50-69%, 6=70-84%, 7=85-95%, 8=96-100.

Numbers in parentheses indicate relative rank in column.