

TABLE 21. 2003 SISKIYOU WINTER WHEAT TEST

Entry	Name	Yield (lbs/acre)	Test Wt (lbs/bu)	1000		Protein (12% mb)	Shatter	Plant Ht (in)	Lodging		Stripe Rust
				Kernel Wt (g)					(soft dough)	(harvest)	
136	YAMHILL	5620 (12)	60.0	45.0	12.28	1.3	41	1.0	1.0	1.5	
397	STEPHENS	6350 (3)	62.0	49.1	12.12	1.5	43	1.0	4.5	1.3	
668	MALCOLM	5680 (11)	62.3	47.0	12.16	2.5	44	1.0	1.8	2.8	
960	ROD	6340 (4)	59.3	41.0	12.12	1.3	40	1.0	3.0	1.0	
967	ROHDE	6640 (1)	64.2	39.3	12.71	1.0	41	1.0	2.5	1.0	
1058	LAMBERT	6540 (2)	62.1	45.1	12.27	2.5	45	1.0	2.8	2.8	
1136	BOUNDARY	6010 (7)	63.0	41.6	11.95	1.3	44	1.0	2.5	5.5	
1198	WEATHERFORD	5830 (9)	60.5	40.9	12.92	1.5	46	1.0	3.3	1.0	
1199	TEMPLE	4830 (17)	63.3	41.2	12.69	1.0	43	3.3	4.8	1.0	
1282	BRUEHL	5280 (15)	59.2	39.3	11.73	1.3	47	4.0	7.3	1.0	
1283	HILLER	5540 (13)	60.4	36.1	12.00	1.3	43	1.0	5.8	1.3	
1284	EDWIN	4400 (19)	62.9	35.0	13.12	1.0	46	7.0	7.8	4.0	
1285	CODA	5880 (8)	62.4	35.4	13.02	1.0	41	2.0	6.0	1.0	
1286	FINLEY	4700 (18)	64.3	41.8	13.14	1.3	51	4.3	5.3	1.0	
1387	GOLDEN SPIKE	6250 (5)	62.3	36.1	12.23	1.3	48	2.3	4.5	2.3	
1388	DELORIS	5240 (16)	64.0	34.6	11.37	1.5	45	1.5	1.5	7.3	
1389	BRUNDAGE 96	6190 (6)	61.3	37.8	12.77	2.0	41	1.0	1.0	3.0	
1390	HUBBARD	5350 (14)	61.6	36.6	12.33	2.8	49	1.0	2.3	5.3	
1413	WINCAL	5730 (10)	64.8	38.5	12.86	1.3	39	1.0	1.0	4.3	
	MEAN	5710	62.1	40.1	12.41	1.5	44	1.9	3.6	2.5	
	CV	13.1	1.0	3.7	-	40.3	3.2	41.9	41.0	21.5	
	LSD (.05)	1060	1.3	3.1	-	0.8	3	1.1	2.1	0.8	

Rating scale for diseases (area of flag-1 leaf affected), lodging and shatter: 1 = 0-3%, 2 = 4-14%, 3 = 15-29%,
4 = 30-49%, 5 = 50-69%, 6 = 70-84%, 7 = 85-95%, 8 = 96-100%.

Numbers in parentheses indicate relative rank in column.