

TABLE 63. 2005 STANISLAUS CEREAL FORAGE STRIP TEST

Entry	Type	Harvest stage	Tons/Acre (70% mb)	Plant Ht (in)			Lodging*	
				12/21	1/6	3/9	1/6	3/9
Cayuse	Oat	Early Head	11.0	18.3	23.0	47.3	L1	L1
Everleaf 114	Oat	Boot	11.2	23.3	28.0	40.0		L1
Kanota	Oat	Flower	12.9	20.7	21.7	47.5		L3
Dirkwin	Wheat	Early Head	11.2	16.0	17.0	35.0		
Super Dirkwin	Wheat	Early Head	11.8	18.3	21.7	39.0		
Forerunner	Triticale	Boot	9.8	12.3	5.0	42.7	L2	
Merlin	Triticale	Boot	9.6	18.0	18.3	41.7		L2
Trical 2700	Triticale	Boot	9.2	18.3	18.7	41.3	L1	

Forage yield differences between entries are not statistically significant

*Lodging: L1= 1 rep lodged; L2= 2 reps lodged; L3 = 3 reps lodged.

Location: Stanislaus Farm Supply Reseach Farm, Modesto

Soil Type: Tujunga loamy sand

Irrigated up: 9/30/04-10/1/04

Preplant fertilizer: broadcast 300 lbs/A ammonium sulfate preplant,

Fertilizer applications: 1-19-05 150 lbs/A ammonium sulfate

Plot size: 12' x 67', commercially swathed, 12' x 20' hand weighed