

Crop Classification	Variety	UC Number	Avg. Yield (lb/ac)	St.Dev., Yield	Trial Coef.Var., Yield	Rank, Yield	Avg. Protein (%)	Rank, Protein	Test Weight (lb/bu)	Thousand Kernel Wt (g)	Plant Height (in)	Lodging, harvest (1-8)
6RSF	UC 960	960	5569	1150	0.21	1	8.0	8	46.4	61.0	32	1
6RSF	UC 933	933	4979	901	0.21	2	8.9	7	45.7	59.5	31	1
6RSF	ISHI	1047	4423	1795	0.21	3	7.4	9	46.1	58.5	34	1
6RSF	UC 603	603	3602	558	0.21	4	12.5	2	44.4	48.5	32	1
6RSN	UC 937	937	3431	1066	0.21	5	11.7	5	44.9	56.5	34	1
6RSF(H)	SCHALLER	1355	3257	-	0.21	6	10.3	6	42.3	51.5	43	1
2RSM	UC TAHOE	1409	2277	-	0.21	7	12.0	4	47.9	51.0	33	1
6RSF	UC 969	969	1915	287	0.21	8	12.4	3	47.6	54.5	33	1
6RSN	TAMALPAIS	1134	1779	92	0.21	9	13.5	1	52.1	53.0	31	1

Rating scale, area of upper leaves affected except for BYDV which is measured as a percentage of plants showing symptoms:

- 1 = 0-3%
- 2 = 4-14%
- 3 = 15-29%
- 4 = 30-49%
- 5 = 50-69%
- 6 = 70-84%
- 7 = 85-95%
- 8 = 95-100%

\*The barley trial at the 2022 Delta location had severe bird damage. Reported values represent varieties and plots for which sufficient grain remained to effectively harvest via small plot combine. Because the degree of bird damage was a local/atypical phenomena, these data are not included in the multi-year, multi-site summaries.